


# “Introducing the Collection 6 MODIS Aerosol products”

## Season III



# Starring: The “Dark Target” Aerosol Team



**Disclaimer:** Lorraine is an renegade superhero. She receives no financial contribution from the MODIS aerosol group, and does not in any way "influence the government on this particular matter on behalf of UMBC".

Telling Rob what to do, on a personal level, is perfectly legal. That's just female-male communication.

When we last saw our  
motley gang of heroes...

(Scenes from Season II)

AeroCenter update: May 2012

# ... They were promising “final” delivery of Collection 6 algorithms

(imagine camera pans into each office)

- Rob was fussing with gas correction
- Leigh was fixing the cloud mask
- Shana was ordering that Version V6.0.11 was the *final* version
- Rich was evaluating DeepBlue/DarkTarget merge
- Lorraine was at UMBC, starting AirPhoton
- Falguni and Pawan had not yet joined the gang.





Now one year later...

(Scenes from Season III)

# ... They are still promising “final” delivery of Collection 6 algorithms

(imagine camera pans into each office)

- **Rob** is answering 40 pages of AMTD comments
- **Leigh** is fixing the coastline mask
- **Shana** is insisting that Version V6.0.27 is the *final* version
- **Rich** is re-evaluating DeepBlue/DarkTarget merge
- **Lorraine** comes to GSFC because she misses the rest of the gang.
- **Falguni** is fussing with gas correction.
- **Pawan** is improving urban aerosol retrieval



# Introducing the Collection

## 6 MODIS aerosol products

Subtitle: The art of the MODIS  
aerosol retrieval

Sub-subtitle: Getting a cat into a carrier


Trying to get C6 into product is like...




Video found on You-Tube  
Note: no harm to cat

# Aerosol retrieval from MODIS

What MODIS observes



Attributed to aerosol (AOD)



There are many different “algorithms” to retrieve aerosol from MODIS

1. Dark Target (“DT” ocean and land; Levy, Mattoo, Munchak, Remer, Tanré, Kaufman)
2. Deep Blue (“DB” desert and beyond; Hsu, Bettenhausen, Sayer,.. )
3. MAIAC (coupled with land surface everywhere; Lyapustin, Wang, Korkin,...)
4. Ocean color/atmospheric correction (McClain, Ahmad, ...)
5. Etc (neural net, model assimilation, statistical, ... )
6. Your own algorithm (many groups around the world)


# Generic MODIS-DT retrieval algorithm

## INPUTS / UPSTREAM PRODUCTS

- MOD02 = MODIS spectral reflectance pixels (at 500 m resolution)
- MOD03 = geo-location: geometry and surface type ( 1 km resolution)
- MOD35 = cloud mask, including IR tests. Also has surface type
- Ancillary data: GDAS 1° ozone, water vapor, surface wind speed

Aggregate into  
N x N boxes

= 20x20 for 10 km





# Last November we submitted Collection 6 overview paper to AMTD

[Levy, R., S. Mattoo, L.A. Munchak, L.A. Remer, A.M. Sayer, and N. Hsu](#) (2013). The Collection 6 MODIS Aerosol Products over Land and Ocean *Atmos. Meas. Tech. Disc.* (6), 159-259


- We thought it would be a service to our users to have paper out when C6 processing began.
- It was a really long paper (100 AMTD pages) describing major updates and effects on the “products”
- It received 40 pages of online criticism and comments
- Then the rug was pulled out from under us.

What happened?

# The world changed (or at least the MODIS-observed view of the world)

## INPUTS / UPSTREAM PRODUCTS

- MOD02 = MODIS spectral reflectance pixels (at 500 m resolution)
- MOD03 = geo-location: geometry and surface type ( 1 km resolution)
- MOD35 = cloud mask, including IR tests. Also has surface type
- Ancillary data: GDAS 1° ozone, water vapor, surface wind speed



- For MOD02: Different decisions about bad/missing pixels, plus calibration
- For MOD03: Definitions of land/water/coastal were changed
- For MOD35: IR test finding high cirrus clouds was changed
- For ozone: Input stream changed from TOAST to GDAS

**“Our” changes for C6 needed to be revised**

Note: that “our” also refers to **ALL** MODIS products (clouds, land surface, etc) <sup>12</sup>


# “Definitions” of land/coastal/water:

## C5: Used MOD35; C6: Use MOD03 (It’s a long story)


This is ARCTAS  
region in  
northern  
Canada.

Land ‘o lakes


MODIS AOD, with no coastline pixels




MODIS AOD, with coastline pixels



improved  
coverage near  
water





C5 MOD03 Land/Sea Mask



Note very little  
“coastal” in C5.

Clean divide  
between Green  
(land) and  
Yellow (Lakes)

C6 MOD03 Land/Sea Mask



Note much  
more “coastal”  
in C6.

Lots of  
shorelines (lime  
colored) pixels  
that would be  
thrown out.

# ... Now they are promising “final” delivery of Collection 6 algorithms

(Shana will allow NO MORE CHANGES!!!!)

## OUTLINE of REST of TALK




- Compensated for upstream changes!!!!
- Better RT (gas corrections, LUTs)
- Cloud mask and pixel decision-making
- “Science” upgrades
- “Quality assurance” upgrades
- DT products (expected C6 versus C5)
- Other stuff: 3 km, DT/DB merge, etc

# Better RT: Gas Corrections

- Gases ( $\text{H}_2\text{O}$ ,  $\text{O}_3$ ,  $\text{CO}_2$ ,  $\text{NO}_2$ , etc... ) absorb radiation,  $T_{\text{gas}} < 1.0$
  - $T_{\text{gas}}$  varies spectrally and function of gas optical depth:
  - $T_{\text{gas}} \approx 1 - \tau_{\text{gas}}$ :
- 
- The C5 assumptions were undocumented (circa 1995)
  - Used modern RT code and found changes
  - Changes for C6 have impacts.

# Gas Corrections (cont.)



- Consistent “bias” of  $\Delta \rho^* \approx 0.001 \rightarrow$  “bias” of AOD of 0.01.
- Bias is **double** for  $Z=60^\circ$ . **Quadruple** for two-way transmission!
- Something really weird is happening at 1.24 μm channel (**big impact on land retrieval**)
- **ACE: We want to know global AOD to  $\pm 0.01$ ? Not when we don't know gas!!!**



# Radiative Transfer updates (applied to C5)


Jul 2008: Aqua

Recalculate “center wavelengths” from MODIS filter functions, and spectral Rayleigh optical depths

Recalculate gas absorption coefficients and optical depths for  $\text{H}_2\text{O}$ ,  $\text{O}_3$ ,  $\text{CO}_2$ ,  $\text{CH}_4$ ,  $\text{O}_2$ , etc


Extend valid solar zenith angle from  $72^\circ$  to  $84^\circ$ .

Pretty big changes, but they are justified.




# Update cloud masking (ocean & land)


MYD04\_L2.A2010227.1705.hdf




Without cloudmask



With new cloudmask



New Pixels



AOD at 550 nm




- Retrieve more “heavy smoke” cases
- Retrieve fewer “thin cirrus” cases
- Correct assignment of QAC due to clouds





# Science: wind speed dependence (ocean)

MYD04\_L2.A2010018.1440.hdf






NCEP winds - 6 m/s winds



- Higher wind speeds → more ocean foam and diffused glitter pattern


- Aerosol LUT now calculated for 4 wind speeds

( $v=2, 6^*, 10, 14$  m/s)



- Retrieved AOD reduced when  $v \geq 6$
- Reduces AOD near  $40^\circ$  glint mask edges and in "Roaring 40s" of southern Oceans.

# Quality Confidence “definitions”



Old (C5): Quality assurance depends on success of a retrieval solution. If aerosol signal is too small, then do not include in daily/monthly statistics.





New (C6) : Sometimes signal is small, but we are confident that it is small. Include in daily/monthly statistics.

- But “major” influence on estimating regional and global mean AOD

# Major bug fix: surface reflectance relationship to NDVI\_swir










Jul 2008

- Coding error from C5 continued to C6
- Surface reflectance dependence on NDVI\_swir was reversed
- New: Reduces super-high bias for semi-arid regions.



These labels were reversed  
Resulting in coding error









# Results from **ALL** changes

Jul 2008: Aqua



Version 6.0.27



NDVI switch

Mostly wind  
speed  
upgrades

Decisions about  
land/sea/coastal

Higher zenith  
angle valid


AOD Difference  
-0.12 -0.08 -0.04 0.00 0.04 0.08 0.12

AOD at 550 nm  
0.0 0.1 0.2 0.3 0.4 0.5 0.6 0.7 0.8

- lower AOD over ocean, but significantly lower over southern oceans
- Higher AOD over most land, but lower in semi-arid regions.
- Changes in decisions to retrieve (see Aral sea).


# Preliminary “validation” of C6

## 6 months testing (on Aqua)



- No dramatic changes, but 10-20% more valid collocations with AERONET
- C6 algorithm has slightly better (68% vs 64%) retrieved over ocean
- C6 will have more of a high bias over land, but slightly better correlation.

## C5: Trends Terra ≠ Aqua




- Over land, **Terra** decreases (-0.04/decade), **Aqua** constant
- **Terra** / **Aqua** divergence is the same everywhere on the globe!
- **Artificial Terra trend traced to degrading of Terra's calibration**



# C6: Impact of New Terra calibration


Jul 2008: Terra



- Main impacts over land
  - Global increase by 0.02 (for this particular month)
- Impact over ocean is small
  - Global increase by 0.004 (for this particular month)

Jack Xiong,  
Junqing Sun and  
MCST

## C6: Impact of new calibration on trend?




- 6 months processed with same dark-target aerosol algorithms
- Terra (T) now “in sync” with Aqua (A) time series
- (Terra-Aqua) offset remains 0.01 (ocean) and 0.015 (land)
- **New calibration → Terra/Aqua divergence removed for C006!**

# What else for C6 Level 2?




## New Diagnostic SDSs

Land\_Ocean\_Quality\_Flag




Topographic\_Altitude\_Land




Surface Height [km]



Land\_Sea\_Flag



Wind\_Speed\_Ncep\_Ocean




Wind Speed [m/s]




# Cloud and Aerosol SDSs

**Aerosol\_Cldmask\_Land\_Ocean**



500 meter resolution cloud mask used in aerosol retrieval. Can be (at times, significantly) different than MOD35

**Distance to Cloud**



Number of pixels between an aerosol retrieval and the closest cloud. Not thoroughly validated yet.


Jan 21 2010 at 13:40 UTC




# The Deep/Dark merge

- **Dark**, **bright**, and **transitional** regions are identified by monthly mean NDVI
- In **Dark** regions, value from dark-target retrieval is used
- In **bright** regions, deep-blue is used
- In **transition** regions, AOD is merged, dependent on QA of retrievals

DeepDark Algorithm Flag




DeepDark 550 nm AOD



C. Hsu, A. Sayer, C. Bettenhausen, S. Mattoo, L. Munchak, R. Kleidman, et al.,



# Monthly mean AOD for Aqua, January 2010






Merging deep blue & dark target produces best global coverage

- Deep blue is land-only; need dark target for oceans
- Deep blue introduces coverage over Australian outback, Sahara desert and Arabian peninsula
- Still no coverage over snow (see: most of Northern Hemisphere).

**MERGE LOOKS REASONABLE, BUT NOT VALIDATED YET!!!! (Kleidman et al)**

# What about C6 Level 3?

## Calculating mean AOD




*Levy et al., TRGS, 2009*

- It depends on the post-processing path
- Many choices for aggregation and weighting
- Accentuate different limitations of original (L2) sampling



# Changes to Level 3: monthly



- Pixel weighting for monthly data is removed
- Replaced by “equal day” weighting (assuming at least 5 observations per day).
- → Increase monthly mean AOD over land, and ocean

# Global → Local






- Nominal resolution (e.g. 10 km) cannot resolve higher scale (e.g. urban, neighborhood) variability
- Air Quality community has been asking (no, *pleading!*) for higher resolution data
- We have been afraid to lose the advantage (pixel statistics) of the 10 km.

# 3 km algorithm

In two AMT-Discussion Papers (in revision for AMT)

- [Munchak, L., R.C. Levy, S. Mattoo, L.A. Remer, B.N. Holben, J.S. Schafer, C.A. Hostetler, and R.A. Ferrare \(2013\). MODIS 3km Aerosol Product: applications over land in an urban/suburban region, Atmos. Chem. Phys. Dis, 6, 1683-1716  
doi:10.5194/amtd-6-1683-2013](#)
- [Remer, L., S. Mattoo, R.C. Levy, and L. Munchak \(2013\). MODIS 3km Aerosol Product: Algorithm and Global Perspective, Atmos. Meas. Tech. Disc., \(6\), 69-112  
doi:10.5194/amtd-6-69-2013](#)
- [J. M. Livingston, J. Redemann, Y. Shinozuka, R. Johnson, P. B. Russell, Q. Zhang, S. Mattoo, L. Remer, R. Levy, L. Munchak, and S. Ramachandran, \(2013\). Comparison of MODIS 3-km and 10-km resolution aerosol optical depth retrievals over land with airborne Sunphotometer measurements during ARCTAS summer 2008, Atmos. Chem. Phys. Dis, \(submitted last week\).](#)

# MODIS 3 km product over suburban (MD) landscape (summer 2010)




- 3 km mirrors 10 km product (pattern and magnitude)
- 3 km introduces **noise**, but also can reduce spatial impact of **outliers**



# MODIS vs DRAGON


## July 21, 2011

AOD (AERONET: DRAGON)





- DRAGON = Many AERONET over Maryland during DISCOVER-AQ experiment, July 2011
- For MODIS plots, QA=3 only
  - Circle Center = AERONET  $\pm 2$  hours
  - Circle Outer = MODIS 5 x 5 box
- 3km (bottom) resolves Baltimore maxima
- Gray is MODIS-Aerosol cloud mask
- 3km also resolves aerosol over Ches. Bay


MYD04\_10km 550 um 2011202 18:30



MYD04\_03km 550 um 2011202 18:30







## MODIS 3 km product (operational for C006)

- Algorithm structurally identical to standard “10 km” retrieval
- Results will be in new files, ***‘MOD04\_3K’***
- Over land and ocean, (but no Deep Blue yet).
- Both MOD04\_L2 and MOD04\_3K will be available


# Summary (1)

- Dark-target algorithm/products are updated for C6, but only “modest” changes (no major science updates).
- **“Modest” is relative...** Many difficult decisions about seemingly small things that actually were significant. (e.g. Gas Correction, Cloud Mask, Quality Assurance etc).
- Upstream changes (L1B and MOD35) meant we had to re-analyze, over and over.
- Not superhero like, but there were mistakes (bugs carried over from C5) that needed re-analysis.
- Dark-target and Deep-blue products are merged, leading to more global coverage

## Summary (2)

- MODIS aerosol retrieval was intended for *global* climate applications
  - Air quality events are sometimes on urban and local scales
  - Aerosol properties change near clouds
  - → The MODIS dark-target team is offering 3 km operational aerosol data.
  - Issues in land/sea/coastal are ACCENTUATED!

# Trying to get C6 into product is like...



Hopefully, there will be no season IV for “Introducing Collection 6 products”



# On towards Collection 7!



- Proper per-pixel error products
- Better urban retrievals
- Use IR information for dust detection
- Etc: It's proposal season!

THANK YOU FOR YOUR ATTENTION !!!